

MATHEMATICS

11

INTERMEDIATE  
PART 1

Bilal Articles

# Chapter 1.

## NUMBER SYSTEMS

# Bilal's Edu & Jobs News

Bilal's Edu & Jobs News on [newsongoogle.com](https://newsongoogle.com/): Navigating the Future of  
Education and Careers

A project of: [www.https://newsongoogle.com/](https://newsongoogle.com/)  
Contact Us: [bilalarticles1@gmail.com](mailto:bilalarticles1@gmail.com)

## Contents

<b>Exercise No.1.1</b> .....	3
<b>Exercise 1.2</b> .....	6
<b>Exercise 1.3</b> .....	10

# Bilal's Edu & Jobs News

Bilal's Edu & Jobs News on [news.google.com](https://news.google.com): Navigating the Future of  
Education and Careers

**1. Rational number**

A number which can be written in the form of

$\frac{p}{q}$  where  $p, q \in \mathbb{Z}$  and  $q \neq 0$  called rational number

Example  $\frac{1}{2}, 4, \frac{3}{4}$

**2. Irrational number**

A Number which cannot be written in the form

of  $\frac{p}{q}$  where  $p, q \in \mathbb{Z}$  and  $q \neq 0$  is called irrational

number. Example  $\sqrt{2}, \sqrt{3}, \sqrt{5}$

**3. Decimal Representation of Rational and Irrational Numbers****1) Terminating decimals:**

A decimal that contains finite number of digits in its decimal part is called Terminating decimal

e.g. 2.02.04, 0.000415

\* Every terminating decimal can be converted to a common fraction.

Every terminating decimal represents a rational number.

**2) Non-Terminating decimals:**

A decimal having infinite number of digits in its decimal part is called non-terminating decimal

e.g., 0.428571...., 0.33333 ...

$\sqrt{2} = 1.414213...$

**There are two types of non-terminating decimals:**

i) Recurring decimals ii) Non-recurring decimals

**i) Recurring decimals**

A recurring or periodic or cyclic decimal is a decimal in which one or more digits repeat indefinitely.

e.g.,  $2.\bar{3} = 2.3333...$  (Rational no.)

$\frac{1}{3} = 0.33333...$  (Rational no.)

$\frac{3}{7} = 0.4285714285714...$  (rational)

Every recurring decimal can be converted to a common fraction.

\* Every recurring decimal represents a rational number.

ii) Non-recurring decimals:

A decimal which neither terminates nor it is recurring is called non-recurring decimal.

e.g.  $\sqrt{2} = 1.414213562...$ ,  $\sqrt{7} = 2.645751...$

Every non-recurring decimal can not be converted to a common fraction.

\* Every non-terminating and non-recurring

decimal represents an irrational number.

**Example 1**

- i.  $0.25 = \frac{25}{100}$  rational no.
- ii.  $\frac{1}{3} = 0.333...$  = (recurring decimal (rational no.))
- iii.  $2.\bar{3} = 2.333...$  Rational no.
- iv.  $0.142857142857 \dots = \frac{1}{7}$  (rational no.)
- v.  $0.01001000100001...$  Non-terminating,

Non-periodic so irrational no.

- vi.  $214.121122111222...$  Irrational no.
- vii.  $1.4142135...$  Is an irrational no.
- viii.  $7.3205080...$  Irrational no.
- ix.  $1.709975947...$  Irrational no.
- x.  $3.141592654...$  Important irrational

number called  $\pi$  (Pi) and  

$$\pi = \frac{\text{circumference of any circle}}{\text{length of its diameter}}$$

**Example 2.**

**Prove that  $\sqrt{2}$  is an irrational number.**

**Solution:**

Suppose  $\sqrt{2}$  is a rational number then  $\sqrt{2} = \frac{p}{q}$  where  $p, q \in \mathbb{Z}$   $\wedge q \neq 0$  if  $HCF(p, q) \neq 1$

then by dividing  $p$  and  $q$  by  $HCF(p, q)$ ,  $\sqrt{2}$  can be reduced as

$\sqrt{2} = \frac{p}{q}$  where  $HCF(a, b) = 1 \rightarrow (1)$

$$\Rightarrow \sqrt{2}b = a$$

$$\Rightarrow 2b^2 = a^2$$

$$\Rightarrow a^2 \text{ is divisible by } 2$$

$$\Rightarrow a \text{ is divisible by } 2 \rightarrow (2)$$

$$\Rightarrow a = 2c \text{ where } c \text{ is an integer}$$

$$\therefore \sqrt{2}b = 2c$$

$$2b^2 = 4c^2$$

$$\Rightarrow b^2 = 2c^2$$

$$\Rightarrow 2b^2 = 4c^2$$

$$\Rightarrow b^2 = 2c^2$$

$$\Rightarrow b \text{ is divisible by } 2 \rightarrow (3)$$

from (2) and (3) 2 is a common factor of  $a$  and  $b$  which contradicts (1)

so  $\sqrt{2}$  is an irrational number.

Prove that  $\sqrt{3}$  is an irrational number.

**Solution:**

Suppose  $\sqrt{3}$  is a rational number then  $\sqrt{3} = \frac{p}{q}$  where  $p, q \in \mathbb{Z} \wedge q \neq 0$  if  $HCF(p, q) = 1$

then by dividing  $p$  and  $q$  by  $HCF(p, q)$ ,  $\sqrt{3}$  can be reduced as

$$\sqrt{3} = \frac{p}{q} \text{ where } HCF(p, q) = 1 \rightarrow (1)$$

$$\Rightarrow \sqrt{3}b = a$$

$$\Rightarrow 3b^2 = a^2$$

$$\Rightarrow a^2 \text{ is divisible by } 3$$

$$\Rightarrow a \text{ is divisible by } 3 \rightarrow (2)$$

$$\Rightarrow a = 3c \text{ where } c \text{ is an integer}$$

$$\therefore \sqrt{3}b = 2c$$

$$3b^2 = 9c^2$$

$$\Rightarrow b^2 = 3c^2$$

$$\Rightarrow b^2 \text{ is divisible by } 3$$

$$\Rightarrow b \text{ is divisible by } 3 \rightarrow (3)$$

from (2) and (3) 3 is a common factor of  $a$  and  $b$  which contradicts (1)

so  $\sqrt{3}$  is an irrational number.

## Properties of real numbers

### Binary operation:-

A binary operation in a set  $A$  is a rule usually denoted by  $*$  that assigns to any pair of elements of  $A$ , taken in a definite order, another element of  $A$ .

\* two binary operations addition and multiplication (.or  $\times$ ) in a set of real numbers ( $\mathbb{R}$ ) are important.

#### 1. Addition laws

##### i) Closure law

$$\forall a, b \in \mathbb{R}, a + b \in \mathbb{R}$$

##### ii) Associative law

$$\forall a, b, c \in \mathbb{R}, a + (b + c) = (a + b) + c$$

##### iii) Additive identity

$$\forall a \in \mathbb{R}, \exists 0 \in \mathbb{R} \text{ such that}$$

$$a + 0 = 0 + a = a \quad 0 \text{ (zero) is called identity element of addition.}$$

##### iv) Additive inverse

$$\forall a \in \mathbb{R}, \exists (-a) \in \mathbb{R} \text{ such that}$$

$$a + (-a) = 0 = (-a) + a$$

##### v) Commutative Law

$$\forall a, b \in \mathbb{R}, a + b = b + a$$

#### 2. Multiplication Laws

##### vi) Closure Law

$$\forall a, b \in \mathbb{R}, a \cdot b \in \mathbb{R}$$

( $a, b$  is usually written as  $ab$ )

##### vii) Associative Law

$$\forall a, b, c \in \mathbb{R}, a \cdot (b \cdot c) = (a \cdot b) \cdot c$$

##### viii) Multiplicative Identity

$$\forall a \in \mathbb{R}, \exists 1 \in \mathbb{R} \text{ such that}$$

$$a \cdot 1 = 1 \cdot a = a \quad 1 \text{ (one) is called identity element of multiplicative.}$$

##### ix) Multiplicative Inverse

$$\forall a \in \mathbb{R}, \exists (a^{-1}) \in \mathbb{R} \text{ such that}$$

$$a \cdot a^{-1} = a^{-1} \cdot a = 1 \quad (a^{-1} \text{ is also written as } \frac{1}{a})$$

$$as \frac{1}{a}$$

##### x) Commutative Law

$$\forall a, b \in \mathbb{R}, ab = ba$$

#### 3. Multiplicative – addition Law

##### xi)

$$\forall a, b, c \in \mathbb{R},$$

$$a(b + c) = ab + ac \text{ left distributive}$$

$$(a + b)c = ac + bc \text{ right distributive}$$

#### 4.

Properties of equality

##### i) Reflexive property

$$\forall a \in \mathbb{R}, a = a$$

## ii) Symmetric Property

$$\forall a, b \in \mathbb{R}, a = b \Rightarrow b = a$$

## iii) Transitive Property

$$\forall a, b, c \in \mathbb{R}, a = b \wedge b = c \Rightarrow a = c$$

## iv) Additive property

$$\forall a, b, c \in \mathbb{R}, a = b \Rightarrow a + c = b + c$$

## v) Multiplicative property

$$\forall a, b, c \in \mathbb{R}, a = b \Rightarrow ac = bc \wedge ca = cb$$

## vi) cancellation property w.r.t addition

$$\forall a, b, c \in \mathbb{R}, a + c = b + c \Rightarrow a = b$$

## vii) Cancellation property w.r.t multiplication

$$\forall a, b, c \in \mathbb{R}, ac = bc \Rightarrow a = b, c \neq 0$$

## 5. Properties of inequalities

## 1) Trichotomy property

$$\forall a, b \in \mathbb{R}, \text{either } a = b \text{ or } a > b \text{ or } a < b$$

## 2) Transitive property

$$\forall a, b, c \in \mathbb{R}$$

$$1. a > b \wedge b > c \Rightarrow a > c$$

$$2. a < b \wedge b < c \Rightarrow a < c$$

## 3) Additive property

$$\forall a, b, c, d \in \mathbb{R}$$

$$a) 1. a > b \Rightarrow a + c > b + c$$

$$2. a < b \Rightarrow a + c < b + c$$

$$b) 1. a > b \wedge c > d \Rightarrow a + c > b + d$$

$$2. a < b \wedge c < d \Rightarrow a + c < b + d$$

## 4) Multiplicative properties

$$a) \forall a, b, c \in \mathbb{R} \text{ and } c > 0$$

$$i) a > b \Rightarrow ac > bc$$

$$ii) a < b \Rightarrow ac < bc$$

$$b) \forall a, b, c \in \mathbb{R} \text{ and } c < 0$$

$$i) a > b \Rightarrow ac < bc$$

$$ii) a < b \Rightarrow ac > bc$$

$$c) \forall a, b, c, d \in$$

$\mathbb{R}$  and  $a, b, c, d$  are all positive

$$1. a > b \wedge c > d \Rightarrow ac > bd$$

$$2. a < b \wedge c < d \Rightarrow ac < bd$$

Note:

\*  $a$  and  $(-a)$  are additive inverse of each other

By def. inverse of  $(-a)$  is  $a$   $-(-a) = a$

\*  $a$  and  $1/a$  are  $x^{-1}$  inverse of each other.

inverse of  $\frac{1}{a}$  is  $a$  (i.e inverse of  $a^{-1}$ )  $a \neq 0$

$$(a^{-1})^{-1} = a \text{ or } \frac{1}{\frac{1}{a}} = a$$

**Exercise No.1.1**

## 1. Which of the following have closure property w.r.t addition and multiplication

i.  $\{0\}$

Solution:- As  $0+0=0 \in \{0\}$ 

$\Rightarrow \{0\}$  has closure property w.r.t addition

As  $0 \times 0 = 0 \in \{0\}$ 

$\Rightarrow \{0\}$  has closure property w.r.t multiplication

ii.  $\{1\}$

Solution:- As  $1+1=2 \notin \{1\}$ 

$\Rightarrow \{1\}$  does not have closure property w.r.t addition

As  $1 \times 1 = 1 \in \{1\}$ 

$\Rightarrow \{1\}$  has closure property w.r.t multiplication

iii.  $\{0, -1\}$

Solution:-

$$0+0=0 \in \{0, -1\}$$

$$0+(-1)=-1 \in \{0, -1\}$$

$$(-1)+0=-1 \in \{0, -1\}$$

$$(-1)+(-1)=-2 \notin \{0, -1\}$$

$\Rightarrow \{0, -1\}$  does not have closure property w.r.t addition

As  $(-1) \times (-1) = 1 \notin \{0, -1\}$ 

$\Rightarrow \{0, -1\}$  has closure property w.r.t multiplication

iv.  $\{1, -1\}$

Solution:-

$$(-1)+(-1)=-2 \notin \{1, -1\}$$

$\Rightarrow \{1, -1\}$  does not have closure property w.r.t addition

Also as

$$1 \times 1 = 1 \in \{1, -1\}$$

$$-1 \times (-1) = 1 \in \{1, -1\}$$

$$(-1) \times 1 = -1 \in \{1, -1\}$$

$$1 \times (-1) = -1 \in \{1, -1\}$$

$\Rightarrow \{1, -1\}$  has closure property w.r.t multiplication.

## 2. Name the property used in the following equations

i.  $4+9=9+4$

Ans:- commutative property w.r.t addition

ii.  $(a+1)+\frac{3}{4} = a+(1+\frac{3}{4})$

Ans:- Associative property w.r.t addition

iii.  $(\sqrt{3} + \sqrt{5}) + \sqrt{7} = \sqrt{3} + (\sqrt{5} + \sqrt{7})$

Ans:- Associative property w.r.t addition

iv.  $100+0=100$

Ans:- additive identity

v.  $1000 \times 1 = 1000$

Ans:- Multiplicative identity

vi.  $4.1 + (-4.1) = 0$

Ans:- Additive inverse

vii.  $a - a = 0$

Ans:- Additive inverse

viii.  $\sqrt{2} \times \sqrt{5} = \sqrt{5} \times \sqrt{2}$

Ans:- commutative property w.r.t multiplication

ix.  $a(b-c) = ab - ac$

Ans:- Left distribution property

x.  $(x-y)z = xz - yz$

Ans:- Right distribution property

xi.  $4 \times (5 \times 8) = (4 \times 5) \times 8$

Ans:- Associative property w.r.t multiplication

xii.  $a(b+c-d) = ab+ac-ad$

Ans:- Left distribution property

3. Name the property used in the following inequalities

i.  $-3 < -2 \Rightarrow 0 < 1$

Ans:- Additive property

ii.  $-5 < -4 \Rightarrow 20 > 16$

Ans:- Multiplicative property

iii.  $1 > -1 \Rightarrow -3 > -5$

Ans:- Additive property

iv.  $a < 0 \Rightarrow -a > 0$

Ans:- Multiplicative property

v.  $a > b \Rightarrow \frac{1}{a} < \frac{1}{b}$

Ans:- Multiplicative property

vi.  $a > b \Rightarrow -a < -b$

Ans:- Multiplicative property

4. Prove the followings rules of addition.

i.  $\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$

$$\begin{aligned} \text{L.H.S.} &= \frac{a}{c} + \frac{b}{c} \\ &= a \times \frac{1}{c} + b \times \frac{1}{c} \\ &= (a+b) \times \frac{1}{c} \\ &= \frac{a+b}{c} = \text{R.H.S} \end{aligned}$$

ii.  $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd}$

$$\begin{aligned} \text{L.H.S.} &= \frac{a}{b} + \frac{c}{d} \\ &= \frac{a}{b} \times 1 + 1 \times \frac{c}{d} \\ &= \frac{a}{b} \times (d \times \frac{1}{d}) + (b \times \frac{1}{b}) \times \frac{c}{d} \\ &= \frac{a}{b} \times \frac{d}{d} + \frac{b}{b} \times \frac{c}{d} \\ &= \frac{ad}{bd} + \frac{bc}{bd} \\ &= ad \times \frac{1}{bd} + bc \times \frac{1}{bd} \\ &= (ad + bc) \times \frac{1}{bd} \\ &= \frac{ad+bc}{bd} = \text{R.H.S} \end{aligned}$$

5. Prove that  $-\frac{7}{12} - \frac{5}{18} = \frac{-21-10}{36}$

$$\begin{aligned} \text{L.H.S} &= -\frac{7}{12} - \frac{5}{18} \\ &= -\frac{7}{12} \times 1 - \frac{5}{18} \times 1 \\ &= -\frac{7}{12} \times (3 \times \frac{1}{3}) - \frac{5}{18} \times (2 \times \frac{1}{2}) \\ &= -\frac{7}{12} \times \frac{3}{3} - \frac{5}{18} \times \frac{2}{2} \\ &= -\frac{21}{36} - \frac{10}{36} \\ &= (-21 - 10) \times \frac{1}{36} \\ &= \frac{-21-10}{36} = \text{R.H.S} \end{aligned}$$

6. Simplify by justify each step.

i.  $\frac{4+16x}{4}$

Solution:-  $= \frac{1}{4} x (4 + 16x)$   
 $= \frac{1}{4} x (4 \times 1 + 4 \times 4x)$  (Multiplicative identity)  
 $= \frac{1}{4} x 4(1 + 4x)$  (Distributive property)  
 $= 1 \times (1 + 4x)$  (Multiplicative inverse)  
 $= (1 + 4x)$  (Multiplicative identity)

ii.  $\frac{\frac{1}{4} + \frac{1}{5}}{\frac{1}{4} - \frac{1}{5}}$

Solution:-  $\frac{\frac{1}{4} \times 1 + \frac{1}{5} \times 1}{\frac{1}{4} \times 1 - \frac{1}{5} \times 1}$   
 $= \frac{\frac{1}{4} \times (\frac{1}{5} \times 5) + \frac{1}{5} \times (\frac{1}{4} \times 4)}{\frac{1}{4} \times (\frac{1}{5} \times 5) - \frac{1}{5} \times (\frac{1}{4} \times 4)}$   
 $= \frac{\frac{1}{4} \times \frac{5}{5} + \frac{1}{5} \times \frac{4}{4}}{\frac{1}{4} \times \frac{5}{5} - \frac{1}{5} \times \frac{4}{4}}$   
 $= \frac{\frac{5}{20} + \frac{4}{20}}{\frac{5}{20} - \frac{4}{20}}$   
 $= \frac{5 \times \frac{1}{20} + 4 \times \frac{1}{20}}{5 \times \frac{1}{20} - 4 \times \frac{1}{20}}$   
 $= \frac{(5 + 4) \times \frac{1}{20}}{(5 - 4) \times \frac{1}{20}}$   
 $= \frac{(5 + 4)}{(5 - 4)}$

iii.  $\frac{\frac{a}{b} + \frac{c}{d}}{\frac{a}{b} - \frac{c}{d}}$

Solution:-  $\frac{\frac{a}{b} \times 1 + \frac{c}{d} \times 1}{\frac{a}{b} \times 1 - \frac{c}{d} \times 1}$  (multiplicative identity)  
 $= \frac{\frac{a}{b} \times (d \times \frac{1}{d}) + \frac{c}{d} \times (b \times \frac{1}{b})}{\frac{a}{b} \times (d \times \frac{1}{d}) - \frac{c}{d} \times (b \times \frac{1}{b})}$  (multiplicative inverse)  
 $= \frac{\frac{ad}{bd} + \frac{bc}{bd}}{\frac{ad}{bd} - \frac{bc}{bd}}$   
 $= \frac{(ad+bc) \times \frac{1}{bd}}{(ad-bc) \times \frac{1}{bd}} \therefore \frac{a}{b} = a \times \frac{1}{b}$   
 $= \frac{ad+bc}{ad-bc}$

iv.  $\frac{\frac{1}{a} + \frac{1}{b}}{1 - \frac{1}{a} \cdot \frac{1}{b}}$

$= \frac{\frac{1}{a} \times 1 + \frac{1}{b} \times 1}{1 - \frac{1}{a} \cdot \frac{1}{b}}$  (multiplicative identity)  
 $= \frac{\frac{1}{a} \times (b \times \frac{1}{b}) + \frac{1}{b} \times (a \times \frac{1}{a})}{1 - \frac{1}{a} \cdot \frac{1}{b}}$  (multiplicative inverse)  
 $= \frac{\frac{1}{a} \times (\frac{b}{b}) + \frac{1}{b} \times (\frac{a}{a})}{1 - \frac{1}{a} \cdot \frac{1}{b}}$

$$= \frac{b+a}{ab - \frac{1}{ab}}$$

$= \frac{b \times \frac{1}{ab} + a \times \frac{1}{ab}}{ab \times \frac{1}{ab} - 1 \cdot \frac{1}{ab}}$  (multiplicative inverse and multiplicative identity)  
 $= \frac{(b+a) \frac{1}{ab}}{(ab-1) \frac{1}{ab}}$  (dist. property)  
 $= \frac{b+a}{ab-1}$  (cancellation law)

**Complex Numbers**

The numbers of the form  $x + iy$  where  $x, y \in R$  and  $i = \sqrt{-1}$  are called complex numbers. here  $x$  is called real part

and  $y$  is called imaginary part of the complex numbers e. g  $3 + 4i, 2 - \frac{5}{7}i$

- Every real number is a complex number with 0 as its imaginary part.

Consider the equation

$$x^2 + 1 = 0$$

$\Rightarrow x^2 = -1$

$\Rightarrow x = \pm \sqrt{-1}$

$\sqrt{-1} \notin R$  for convenience call it imaginary number and denote it by  $i$  (read  $i$  as *iota*)

**Power of  $i$**

$i^2 = -1$  by def.

$i^3 = i^2 \cdot i = (-1)i = -i$

$i^4 = i^2 \cdot i^2 = (-1)(-1) = 1$

$i^{13} = i^{12} \cdot i = (i^2)^6 \cdot i = (-1)^6 \cdot i$

$(1)i = i$

$i^6 = (i^2)^3 = (-1)^3 = -1$

thus any power of  $i$  must be equal to  $i, -i, 1,$  and  $-1$

**Operation on complex numbers**

1)  $a + bi = c + di$

$\Rightarrow a = c \wedge b = d$

2) Addition

$a + bi + (c + di) = (a + c) + (b + d)i$

3)  $k(a + bi) = ka + kbi$

4)  $(a + bi) - (c + di) = (a - c) + (b - d)i$

5)  $(a + bi) \cdot (c + di) = ac + adi + bci + bdi^2 = (ac - bd) + (ad + bc)i$

**Conjugate complex Numbers:**

for  $z = a + ib$  then its conjugate is denoted by  $\bar{z}$  and is defined as

$$\bar{z} = \overline{a + ib} = a - ib$$

- A real number is self –conjugate.

Complex Numbers of Ordered pairs of Real Numbers.

$$i) (a, b) = (c, d) \Leftrightarrow a = c \wedge b = d$$

$$ii) (a, b) + (c, d) = (a + c, b + d)$$

$$iii) \text{ if } k \text{ is any real number, then}$$

$$k(a, b) = (ka, kb)$$

$$iv) (a, b)(c, d) = (ac - bd, ad + bc)$$

$$v) (a, b) - (c, d) = (a - c, b - d)$$

Properties of the fundamental operation on complex Numbers.

i) the additive identity in  $C$  is  $(0,0)$

ii) every complex number  $(a, b)$  has the additive inverse  $(-a, -b)$

$$i.e (a, b) + (-a, -b) = (0,0)$$

iii) the multiplicative identity is  $(1,0)$

$$i.e (a, b)(1,0) = (a.1 - b.0, b.1 + a.0)$$

iv) every non zero complex number

i.e number not equal to  $(0,0)$  has a multiplicative inverse.

Q.

Prove that the multiplicative inverse of  $(a, b)$

$$\text{Is} \left( \frac{a}{a^2+b^2}, -\frac{b}{a^2+b^2} \right)$$

Proof:

let  $Z = (a, b)$  or  $Z = a + ib$

$$Z^{-1} = \frac{1}{a + ib} = \frac{1}{a + ib} \times \frac{a - bi}{a - bi}$$

$$Z^{-1} = \frac{a - bi}{(a + ib)(a - ib)} = \frac{a - bi}{a^2 - i^2 b^2}$$

$$Z^{-1} = \frac{a - bi}{a^2 - (-1)b^2} = \frac{a - bi}{a^2 + b^2}$$

$$Z^{-1} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i$$

Hence

$$Z^{-1} = \left( \frac{a}{a^2 + b^2}, -\frac{b}{a^2 + b^2} \right)$$

v)  $(a, b)[(c, d) \pm (e, f)]$

$$= (a, b)(c, d) \pm (a, b)(e, f)$$

A special subset of  $C$  For all  $(a, 0), (b, 0) \in C$

$$(a, 0) + (b, 0) = (a + b, 0)$$

$$(a, 0) \cdot (b, 0) = (ac, 0)$$

$$K(a, 0) = (ka, 0)$$

Multiplicative inverse of  $(a, 0)$  is

$$\left( \frac{1}{a}, 0 \right) \text{ provide } a \neq 0$$

### Exercise 1.2

#### 1. Verify the addition properties of complex Number

i. Closure property

For  $(a, b), (c, d) \in C$

$$(a, b) + (c, d) = (a + c, b + d) \in C$$

ii. Associative property

For  $(a, b), (c, d), (e, f) \in C$

$$= [(a, b) + (c, d)] + (e, f)$$

$$= (a + c, b + d) + (e, f)$$

$$= [(a + c) + e, (b + d) + f]$$

$$= [a + (c + e), b + (d + f)]$$

$\therefore$  ' + ' is associative in  $R$

$$= (a, b) + (c + e, d + f)$$

$$= (a, b) + [(c, d) + (e, f)]$$

iii. Additive identity

$\forall (a, b) \in C$  there is  $(0,0) \in C$

Such that  $(a, b) + (0,0)$

$$= (a + 0, b + 0) = (a, b)$$

iv. Additive inverse

$\forall (a, b) \in C$  there is  $(-a, -b) \in C$

Such that  $(a, b) + (-a, -b)$

$$= (a - a, b - b) = (0,0)$$

v. Commutative property

$\forall (a, b), (c, d) \in C$

$$= (a, b) + (c, d)$$

$$= (a + c) + (b + d)$$

$$= (c + a) + (d + b)$$

$$= (c, d) + (a, b)$$

Q2. Verify the multiplication properties of the complex numbers.

Solution:

1. Close w.r.t "x"

$(a + ib), (c + id) \in C$  then

$$(a + ib)(c + id) = ac + iad + ibc + i^2bd$$

$$= ac + i(ad + bc) - bd$$

$$= (ac - bd) + i(ad + bc) \in C$$

2. Associative w.r.t "x"

$$\begin{aligned} &(a + ib), (c + id), (e + if) \in C \\ &[(a + ib)(c + id)](e + if) \\ &= [(ac - bd) + i(bc + ad)](e + if) \\ &= [e(ac - bd) - f(bc + ad) \\ &\quad + i[f(ac - bd) + e(bc + ad)]] \\ &= [eac - ebd - fbc - fad] \\ &\quad + i[fac - fbd + ebc + ead] \\ &= [a(ec - df) - b(df + de)] \\ &\quad + i[a(cf + de) + b(ec - df)] \\ &= (a + ib)[(ec - df) + i(cf + de)] \\ &= (a + ib)[(c + id)(e + f)] \end{aligned}$$

iii) Identity

$$\begin{aligned} &(a + ib), (1 + i0) \in C \text{ then} \\ &(a + ib)(a + i0) = a + 0 + ib + 0 \\ &a + ib \in C \end{aligned}$$

iv) Inverse

$$(a + ib), \left(\frac{a}{a^2 + b^2}, \frac{ib}{a^2 + b^2}\right) \in C$$

Then

$$\begin{aligned} &= (a + ib) \left(\frac{a}{a^2 + b^2} - \frac{ib}{a^2 + b^2}\right) \\ &= (a + ib) \left(\frac{a - ib}{a^2 + b^2}\right) \\ &= \frac{a^2 - (ib)^2}{a^2 + b^2} = \frac{a^2 + b^2}{a^2 + b^2} = 1 = 1 + i0 \end{aligned}$$

v) Commutative

$$\begin{aligned} &(a + ib), (c + id) \in C \\ &(a + ib)(c + id) \\ &= (ac - bd) + i(ad + bc) \\ &= ca - db + i(ad + cb) \end{aligned}$$

$$= (c + id)(a + ib)$$

Q.3 verify the distributive law of complex numbers

$$\begin{aligned} &(a, b)[(c, d) + (e, f)] \\ &= (a, b)(c, d) + (a, b)(e, f) \end{aligned}$$

Solution :

L.H.S

$$\begin{aligned} &(a, b)[(c, d) + (e, f)] \\ &= (a, b)(c + e, d + f) \\ &= (a(c + e) - b(d + f), a(d + f) + b(c + e)) \\ &= (ac + ae - bd - bf, ad + af + bc + ba) \end{aligned}$$

R.H.S

$$\begin{aligned} &= (a, b)(c, d) + (a, b)(e, f) \\ &= (ac - bd, ad + bc) + (ae - bf, af + bc) \\ &= (ac + ae - bd - bf, ad + af + bc + ba) \end{aligned}$$

Hence proved.

2. Simplify the following.

i.  $i^9$

Solution:-  $(i^2)^4 \cdot i$   
 $= (-1)^4 \cdot i$

$= 1 \cdot i$   
 $= i$

ii.  $i^{14}$

Solution:-  $(i^2)^7$   
 $= (-1)^7$   
 $= -1$

iii.  $(-i)^{19}$

Solution:-  $-(i^2)^9 \cdot i$   
 $= -(-1)^9 \cdot i$   
 $= -(-1) \cdot i$   
 $= 1 \cdot i$   
 $= i$

iv.  $(-1)^{-\frac{21}{2}}$

Solution:

$$\begin{aligned} \frac{1}{(-1)^{\frac{21}{2}}} &= \frac{1}{\left[(-1)^{\frac{1}{2}}\right]^{21}} \\ &= \frac{1}{i^{21}} = \frac{1}{(i^2)^{10} \cdot i} \\ &= \frac{1}{(-1)^{10} \cdot i} = \frac{1}{1 \cdot i} \\ &= \frac{1}{i} \times \frac{i}{i} \\ &= \frac{i}{i^2} = \frac{i}{(-1)} \\ &= -i \end{aligned}$$

Q5.

Write into the term of  $i$

i.  $\sqrt{-1}b$

Solution:-  $ib$

ii.  $\sqrt{-5}$

solution:-  $\sqrt{-1} \times \sqrt{5}$   
 $= \sqrt{-1} \sqrt{5} = \sqrt{5} i$

iii.  $\sqrt{-\frac{16}{25}}$

Solution:-  $\sqrt{-1} \times \sqrt{\frac{16}{25}}$   
 $= \sqrt{-1} \times \sqrt{\frac{16}{25}}$   
 $= \frac{4}{5} i$

iv.  $\sqrt{\frac{1}{-4}}$

Solution:-  $\sqrt{-1} \times \sqrt{\frac{1}{4}}$   
 $= \sqrt{-1} \times \sqrt{\frac{1}{4}} = i \times \frac{1}{2}$

$$= \frac{i}{2} \text{ Ans.}$$

**Q6.**Solve  $(7,9)+(3,-5)$ **Solution:-**

$$\begin{aligned} & (7,9) + (3,-5) \\ &= (7+3, 9-5) \\ &= (10,4) \\ &= \frac{-15 + 40i + 12i - 32i^2}{9 - 64i^2} \\ &= \frac{-15 + 52i + 32}{9 + 64} \quad \therefore i^2 = -1 \\ &\frac{17 + 52i}{73} = \frac{17}{73} + \frac{52}{73}i = \left(\frac{17}{73}, \frac{52}{73}\right) \end{aligned}$$

**Q7.**Solve  $(8,-5)-(-7,4)$ **Solution:-**  $(8+7,-5-4)$ 

$$=(15,-9)$$

**Q8.Solve**  $(2,6) \cdot (3,7)$ **Solution:**

$$\begin{aligned} & (2 + 6i) \cdot (3 + 7i) \\ &= (2 \cdot 3 + 2 \cdot 7i + 6i \cdot 3 + 6i \cdot 7i) \\ &= (6 + 14i + 18i + 42i^2) \\ &= (6 + 42(-1) + 32i) \\ &= (6 - 42 + 32i) \\ &= (-36 + 32i) = (-36, 32) \end{aligned}$$

**Q9.**

$$(5, -4)(-3, -2)$$

**Solution:**

$$\begin{aligned} & (5, -4)(-3, -2) \\ &= (5(-3) - (-4)(-2), 5(-2) + (-4)(-3)) \\ &= (-15 - 8, -10 + 12) = (-23, 2) \end{aligned}$$

**Q10.** $(0,3)(0,5)$ **Solution:** $(0,3)(0,5)$ 

$$\begin{aligned} &= (0.3 - 3.5, 0.5 + 3.0) = (0.15, 0 + 0) \\ &= (-15, 0) \end{aligned}$$

**Q.11****Solve**  $(2,6) \div (3,7)$ **Solution:-**

$$\begin{aligned} & \frac{(2,6)}{(3,7)} = \frac{2+6i}{3+7i} \\ &= \frac{2+6i}{3+7i} \times \frac{3-7i}{3-7i} \\ &= \frac{(2+6i)(3-7i)}{(3)^2 - (7i)^2} \\ &= \frac{(2 \cdot 3 - 2 \cdot 7i + 6i \cdot 3 - 6i \cdot 7i)}{9 - 49(i)^2} \\ &= \frac{6 - 14i + 18i - 42i^2}{9 - 49(-1)} \end{aligned}$$

$$\begin{aligned} &= \frac{6 + 4i - 42(-1)}{9 + 49} = \frac{6 + 42 + 32i}{58} \\ &= \frac{48 + 4i}{58} = \left(\frac{24}{29}, \frac{2}{32}\right) \end{aligned}$$

**Q.12****Solution**

$$\begin{aligned} & (5, -4) \div (-3, -8) \\ &= \frac{5, -4}{(-3, -8)} = \frac{5 - 4i}{-3 - 8i} \\ &= \frac{5 - 4i}{-3 - 8i} \times \frac{(-3 + 8i)}{(-3 + 8i)} \\ &= \frac{5(-3) + (5)(8i) + (-4i)(-3) + (-4i)(8i)}{(-3)^2 - (8i)^2} \end{aligned}$$

**Q.13****Prove that sum as well as product of two conjugate complex number is real.****Solution:-**

let two conjugate complex number be

 $Z = a + ib$  and  $\bar{z} = a - ib$  where  $a, b \in \mathbb{R}$ 

$$\text{Sum} = z + \bar{z} = a + ib + a - ib$$

$$= 2a \in \mathbb{R} \quad \therefore a \in \mathbb{R}$$

$$\text{Product} = z \cdot \bar{z}$$

$$= (a + ib) \cdot (a - ib)$$

$$= (a^2 - i^2 b^2)$$

$$= a^2 - (-1)b^2$$

$$= a^2 + b^2 \in \mathbb{R} \quad \therefore x, b \in \mathbb{R}$$

**Q14.****Find the multiplicative inverse of the following**i.  $(-4,7)$ **Solution:-** let  $z = (-4,7)$ Multiplicative inverse of  $z = \frac{1}{z}$ 

$$= \frac{1}{(-4,7)} = \frac{1}{-4 + 7i}$$

$$= \frac{1}{-4+7i} = \frac{1}{-4+7i} \times \frac{-4-7i}{-4-7i}$$

$$= \frac{-4-7i}{(-4)^2 - (7i)^2} = \frac{-4-7i}{16 - 49(i)^2}$$

$$= \frac{-4-7i}{16+49} = \frac{-4-7i}{65}$$

$$= \frac{-4-7i}{65} = \frac{-4}{65} - \frac{7}{65}i$$

$$= \left(-\frac{4}{65}, -\frac{7}{65}\right)$$

ii.  $(\sqrt{2}, \sqrt{5})$ **Solution:-** let  $z = (\sqrt{2}, -\sqrt{5})$ Multiplicative inverse of  $z = \frac{1}{z}$ 

$$= \frac{1}{(\sqrt{2}, -\sqrt{5})} = \frac{1}{\sqrt{2} - \sqrt{5}i}$$

$$= \frac{1}{\sqrt{2} - \sqrt{5}i} \times \frac{\sqrt{2} + \sqrt{5}i}{\sqrt{2} + \sqrt{5}i}$$

$$= \frac{\sqrt{2} + \sqrt{5}i}{(\sqrt{2})^2 - (\sqrt{5}i)^2}$$

$$= \frac{\sqrt{2} + \sqrt{5}i}{2 - 5i^2}$$

$$= \frac{\sqrt{2} + \sqrt{5}i}{2 - 5(-1)} = \frac{\sqrt{2} + \sqrt{5}i}{2 + 5}$$

$$= \frac{\sqrt{2} + \sqrt{5}i}{7} = \left(\frac{\sqrt{2}}{7}, \frac{\sqrt{5}}{7}\right)$$

iii. (1,0)

Solution:

$$\text{let } z = (1, 0)$$

Multiplicative inverse of  $z = \frac{1}{z}$

$$\frac{1}{(1,0)} = \frac{1}{1+0i}$$

$$= \frac{1}{1+0} = \frac{1}{1} = 1$$

$$= (1,0)$$

Q15.

Factorize the following.

i.  $a^2 + 4b^2$

Solution:-  $a^2 - (-1)4b^2$

$$= a^2 - (i)^2 2^2 b^2$$

$$= (a)^2 - (i2b)^2$$

$$= (a + i2b)(a - i2b)$$

$$= (a + 2bi)(a - 2bi)$$

ii.  $9a^2 + 16b^2$

Solution:-  $9a^2 - (-1)16b^2$

$$= (3a)^2 - (i)^2 4^2 b^2$$

$$= (3a)^2 - (i4b)^2$$

$$= (3a + i4b)(3a - i4b)$$

$$= (3a + 4bi)(3a - 4bi)$$

iii.  $3x^2 + 3y^2$

Solution:-  $3x^2 - (-1)3y^2$

$$= 3(x)^2 - (i)^2 3y^2$$

$$= 3[(x)^2 - (iy)^2]$$

$$= 3[(x + iy)(x - iy)]$$

$$= 3(x + iy)(x - iy)$$

Q17. Separate real and imaginary part.

i.

$$\frac{2 - 7i}{4 + 5i}$$

Solution:-  $\frac{2-7i}{4+5i} \times \frac{4-5i}{4-5i}$

$$= \frac{8-10i-28i+35i^2}{(4)^2 - (5i)^2}$$

$$= \frac{8-38i+35(-1)}{16-25i^2}$$

$$= \frac{8-38i-35}{16-25(-1)}$$

$$= \frac{-27-38i}{16+25} = \frac{-27-38i}{41}$$

$$= \frac{-27}{41} - \frac{38}{41}i$$

$$\frac{(-2 + 3i)^2}{1 + i}$$

Solution:-  $\frac{(-2)^2 + (3i)^2 + 2(-2)(3i)}{1+i}$

$$= \frac{4+9i^2-12i}{1+i} = \frac{4+9(-1)-12i}{1+i}$$

$$= \frac{4-9-12i}{1+i} = \frac{-5-12i}{1+i}$$

$$= \frac{-5-12i}{1+i} \times \frac{1-i}{1-i}$$

$$= \frac{-5+5i-12i+12i^2}{(1)^2 - (i)^2}$$

$$= \frac{-5+5i-12i+12(-1)}{1-(-1)}$$

$$= \frac{-17-7i}{2} = \frac{-17}{2} - \frac{7}{2}i$$

iii.  $\frac{i}{1+i}$

Solution:-  $\frac{i}{1+i} \times \frac{1-i}{1-i}$

$$= \frac{i-i^2}{1^2-i^2} = \frac{i-(-1)}{1-(-1)}$$

$$= \frac{i+1}{1+1} = \frac{i+1}{2}$$

$$= \frac{1}{2} + \frac{i}{2}$$

**The Real Line**

The set of real numbers are represented by a straight line  $XOX'$  as shown.



**The Real Plane /coordinate plane**

The plane made by two mutually perpendicular lines is called coordinate plane. Let us draw two mutually  $\perp$

lines  $XX'$  and  $YY'$  such as  $O$  be their

Point of intersection. The lines

$XX'$  and  $YY'$  are together coordinates axes. The

common point  $O$  is called origin or initial point.

$XOX'$  is called X-axis, which is horizontal line and

$YOY'$  is called Y-axis. Which is vertical line. Thus

the plane made by both x-axis and y-axis is called xy-plane or real plane.


If  $(a,b)$  are called coordinates of a point  $p$  then  $a$

is called x-coordinate or abscissa of point  $p$  and  $b$

is called y-coordinate are ordinate of point  $p$ . the

coordinate plane into four equal parts, called


quadrants.



**Geometrical representation of complex Numbers**

Geometrical representation of complex Number  $a + ib$  is represented by a point P(a,b) on the coordinate plane. When we represent a complex number on coordinate plane, then the coordinate plane is called complex plane or Z plane.


- In the representation the x-axis is called real axis and y-axis is called imaginary axis.



**Argand diagram:**

Figure representing one or more complex numbers on complex plane is called Argand diagram.

Modulus of the Complex Number  $a + ib$



In Cartesian plane distance of A(a,b) from origin O(0,0)

$$|OP| = \sqrt{(a - 0)^2 + (b - 0)^2} = \sqrt{a^2 + b^2}$$

Draw  $\perp AM$  from A on x-axis then  $|OM| = a$

$|AM| = b$  and  $|OA| = z$  by pathagoras

therom on  $\triangle AOM$

$$|OA|^2 = |OM|^2 + |AM|^2$$

$$Z^2 = a^2 + b^2$$

$$|Z| = \sqrt{a^2 + b^2}$$

Thus the modulus of a complex number from the origin.

**Exercise 1.3**

**Question Find multiplicative inverse of each of the following numbers.**

i.  $-3i$

Solution:-let  $z = -3i$  then its multiplicative inverse is  $\frac{1}{z}$

$$\begin{aligned} \frac{1}{z} &= \frac{1}{-3i} \times \frac{i}{i} \\ &= \frac{i}{-3(i)^2} = \frac{i}{-3(-1)} \\ &= \frac{i}{3} \end{aligned}$$

ii.  $1-2i$

Solution:-let  $z = 1-2i$  then its multiplicative inverse is  $\frac{1}{z}$

$$\begin{aligned} \frac{1}{z} &= \frac{1}{1-2i} \\ &= \frac{1}{1-2i} \times \frac{1+2i}{1+2i} \\ &= \frac{1+2i}{(1)^2 - (2i)^2} \\ &= \frac{1+2i}{1-4(i)^2} = \frac{1+2i}{1-4(-1)} \\ &= \frac{1+2i}{1+4} = \frac{1+2i}{5} \\ &= \frac{1}{5} + \frac{2}{5}i \end{aligned}$$

iii.  $-3 - 5i$

Solution:

let  $Z = -3 - 5i = (-3, -5)$

$\therefore$  for  $Z = (a, b)$

$$\Rightarrow z^{-1} = \left( \frac{a}{a^2+b^2}, -\frac{b}{a^2+b^2} \right)$$

so for  $Z = (-3, -5)$

$$\Rightarrow z^{-1} = \left( -\frac{3}{(-3)^2+(-5)^2}, -\frac{-5}{(-3)^2+(-5)^2} \right)$$

$$= \left( \frac{-3}{9+25}, \frac{5}{9+25} \right) = \left( \frac{-3}{34}, \frac{5}{34} \right)$$

$$= \frac{-3}{34} + \frac{5}{34}i$$

iv. (1,2)

Solution:

$$\text{let } Z = (1,2)$$

$$\therefore \text{for } Z = (a, b)$$

$$z^{-1} = \left( \frac{a}{a^2 + b^2}, -\frac{b}{a^2 + b^2} \right)$$

For Z=(1,2)

$$\Rightarrow z^{-1} = \left( \frac{1}{(1)^2 + (2)^2}, \frac{-2}{(1)^2 + (2)^2} \right)$$

$$= \left( \frac{1}{1+4}, \frac{-2}{1+4} \right) = \left( \frac{1}{5}, \frac{-2}{5} \right) = \frac{1}{5} + \frac{-2}{5}i$$

**Question No.3**

Simplify

i.  $i^{101}$

solution:-  $i^{100} \cdot i$

$= (i^2)^{50} \cdot i$

$= (-1)^{50} \cdot i$

$= 1 \cdot i = i$

ii.  $(-ai)^4$

solution:-  $(-a)^4 i^4$

$= a^4 \cdot (i^2)^2$

$= a^4 \cdot (-1)^2$

$= a^4 \cdot 1 = a^4$

iii.  $i^{-3}$

solution:-  $\frac{1}{i^3}$

$= \frac{1}{i^2 \cdot i}$

$= \frac{1}{(-1) \cdot i}$

$= \frac{1}{(-1) \cdot i} = \frac{-1}{i}$

$= \frac{-1}{i} \times \frac{i}{i}$

$= \frac{-i}{i^2} = \frac{-i}{(-1)}$

$= i$

iv.  $i^{-10}$

solution:-  $\frac{1}{i^{10}}$

$= \frac{1}{(i^2)^5}$

$= \frac{1}{(-1)^5}$

$= \frac{1}{-1} = -1$

**Question No.4 Prove that  $\bar{z} = z$  iff  $z$  is real.**Proof :- let  $z = a - bi$  and  $\bar{z} = a - bi$ 

Let  $\bar{z} = z$

Then  $a - bi = a + bi$

$\Rightarrow -bi = bi$

$\Rightarrow bi + bi = 0$

$\Rightarrow 2bi = 0$

$\Rightarrow b = 0$  so  $z = a$  (real)

Conversely let  $z$  be a real number that is  $z = a$ ,  $a \in \mathbb{R}$ Then  $\bar{z} = \bar{a} = a$  because 'a' is a real number.

Hence  $\bar{z} = z$

**Question No.5****Simplify by expressing in the form of  $a + bi$** 

i.  $5 + 2\sqrt{-4}$

Solution:-  $5 + 2\sqrt{4}\sqrt{-1}$

$= 5 + 2(2)i = 5 + 4i$

ii.  $(2 + \sqrt{-3})(3 + \sqrt{-3})$

Solution:-  $(2 + \sqrt{-3})(3 + \sqrt{-3})$

$= (6 + 2\sqrt{-3} + 3\sqrt{-3} + (\sqrt{-3})^2)$

$= (6 + 5\sqrt{-3} + (-3))$

$= (6 + 5\sqrt{3}i - 3)$

$= (3 + 5\sqrt{3}i)$

iii.  $\frac{2}{\sqrt{5} + \sqrt{-8}}$

Solution:-  $\frac{2}{\sqrt{5} + \sqrt{-8}}$

$= \frac{2}{\sqrt{5} - \sqrt{8}i} \times \frac{\sqrt{5} + \sqrt{8}i}{\sqrt{5} + \sqrt{8}i}$

$= \frac{2(\sqrt{5} - \sqrt{8}i)}{(\sqrt{5})^2 - (\sqrt{8}i)^2}$

$= \frac{2(\sqrt{5} - \sqrt{8}i)}{5 - 8(i)^2} = \frac{2(\sqrt{5} - \sqrt{8}i)}{5 - 8(-1)}$

$= \frac{2(\sqrt{5} - \sqrt{8}i)}{5 + 8} = \frac{2(\sqrt{5} - \sqrt{8}i)}{13}$

$= \frac{2\sqrt{5}}{13} - \frac{2\sqrt{8}}{13}$

iv.  $\frac{3}{\sqrt{6} - \sqrt{-12}}$

Solution:-  $\frac{3}{\sqrt{6} - \sqrt{-12}}$

$= \frac{3}{\sqrt{6} - \sqrt{-12}} \times \frac{\sqrt{6} + \sqrt{-12}}{\sqrt{6} + \sqrt{-12}}$

$= \frac{3(\sqrt{6} + \sqrt{-12})}{(\sqrt{6})^2 - (\sqrt{12}i)^2}$

$= \frac{3(\sqrt{6} + \sqrt{-12})}{6 - 12(i)^2} = \frac{2(\sqrt{5} + \sqrt{8}i)}{6 - 12(-1)}$

$= \frac{3\sqrt{6} + \sqrt{12}i}{6 + 12} = \frac{3(\sqrt{6} + \sqrt{12}i)}{18}$

$= \frac{3\sqrt{6}}{18} + \frac{3.2\sqrt{3}i}{18}$

$$\begin{aligned} &= \frac{\sqrt{6}}{6} + \frac{2\sqrt{3}i}{6} \\ &= \frac{1}{\sqrt{6}} + \frac{\sqrt{3}i}{3} \\ &= \frac{1}{\sqrt{6}} + \frac{1}{\sqrt{3}}i \end{aligned}$$

**Question No.6 Show that**

- i.  $\bar{z}^2 + z^2$  is a real number  
Solution:- let  $z=a+ib$  then  
 $\bar{z}=a-ib$   
 $\bar{z}^2 + z^2 = (a+ib)^2 + (a-ib)^2$   
 $= a^2 + b^2i^2 + 2(a)(ib) + a^2 + b^2i^2 - 2(a)(ib)$   
 $= 2a^2 + 2b^2(-1)$   
 $= 2a^2 - 2b^2$  which is real.
- ii.  $\bar{z}^2 - z^2$  is a imaginary number  
Solution:- let  $z=a+ib$  then  
 $\bar{z}=a-ib$   
 $\bar{z}^2 - z^2 = (a+ib)^2 - (a-ib)^2$   
 $= a^2 + b^2i^2 + 2(a)(ib) - a^2 - b^2i^2 + 2(a)(ib)$   
 $= 4abi$   
 $= 2a^2 - 2b^2$  Which is imaginary number.

**Question No.7 Simplify**

- i.  $(-\frac{1}{2} + \frac{\sqrt{3}}{2}i)^3$   
solution:-  $(\frac{-1+\sqrt{3}i}{2})^3$   
 $= \frac{-1+\sqrt{3}i}{2}$   
 $= (\omega)^3 \quad \therefore \omega^3 = 1$   
 $= 1$
- ii.  $(-\frac{1}{2} - \frac{\sqrt{3}}{2}i)^3$   
solution:-  $(\frac{-1-\sqrt{3}i}{2})^3 \quad \therefore \omega^2$   
 $= \frac{-1-\sqrt{3}i}{2}$   
 $= (\omega^2)^3$   
 $= (\omega^3)^2$   
 $= (1)^2 \quad \therefore \omega^3 = 1$   
 $= 1$
- iii.  $(-\frac{1}{2} - \frac{\sqrt{3}}{2}i)^{-2} (-\frac{1}{2} - \frac{\sqrt{3}}{2}i)$   
Solution:-  $(-\frac{1}{2} - \frac{\sqrt{3}}{2}i)^{-2} (-\frac{1}{2} - \frac{\sqrt{3}}{2}i)$   
 $\frac{\sqrt{3}}{2}i$   
 $= (-\frac{1}{2} - \frac{\sqrt{3}}{2}i)^{-2+1}$   
 $= a^3 + b^3(-1).i + 3a^2bi + 3ab^2(-1)$

$$= \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{-1}$$

$$= \frac{1}{-\frac{1}{2} - \frac{\sqrt{3}}{2}i} \times \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}i}{-\frac{1}{2} + \frac{\sqrt{3}}{2}i}$$

$$= \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}i}{(\frac{1}{2})^2 - (\frac{\sqrt{3}}{2}i)^2}$$

$$= \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}i}{\frac{1}{4} - \frac{3}{4}(-1)}$$

$$= \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}i}{\frac{1}{4} + \frac{3}{4}}$$

$$= \frac{-1 + \sqrt{3}i}{2}$$

- iv.  $(a+bi)^2$   
solution:-  $(a)^2 + (bi)^2 + 2abi$   
 $= a^2 + b^2(i)^2 + 2abi$   
 $= a^2 + b^2(-1) + 2abi$   
 $= a^2 - b^2 + 2abi$

- v.  $(a+bi)^{-2}$   
solution:-  $(a+bi)^{-2}$   
 $= \frac{1}{(a+bi)^2}$   
 $= \frac{1}{(a)^2 + (bi)^2 + 2abi}$   
 $= \frac{1}{a^2 + b^2(i)^2 + 2abi}$   
 $= \frac{1}{a^2 + b^2(-1) + 2abi}$   
 $= \frac{1}{(a^2 - b^2) + 2abi}$   
 $\times \frac{(a^2 - b^2) - 2abi}{(a^2 - b^2) - 2abi}$   
 $= \frac{(a^2 - b^2) - 2abi}{(a^2 - b^2)^2 - 4a^2b^2i^2}$   
 $= \frac{(a^2 - b^2) - 2abi}{a^4 + b^4 - 2a^2b^2 - 4a^2b^2(-1)}$   
 $= \frac{(a^2 - b^2) - 2abi}{(a^2 - b^2) - 2abi}$   
 $= \frac{a^4 + b^4 - 2a^2b^2 + 4a^2b^2}{(a^2 - b^2) - 2abi}$   
 $= \frac{(a^2 + b^2)^2}{(a^2 - b^2) - 2abi}$   
 $= \frac{(a^2 - b^2) - 2abi}{(a^2 + b^2)^2} - \frac{2ab}{(a^2 + b^2)^2}i$

- vi.  $(a+bi)^3$   
Solution:-  $(a+bi)^3$   
 $= (a)^3 + (bi)^3 + 3(a)(bi)(a+bi)$   
 $= a^3 + b^3(i)^3 + 3abi(a+bi)$   
 $= a^3 + b^3(i)^2.i + 3a^2bi + 3ab^2(i)^2$

www.notes.pk.com

$$=a^3 - b^3i + 3a^2bi - 3ab^2$$

vii.  $(a - bi)^3$

$$\begin{aligned} \text{Solution:- } & (a - bi)^3 \\ & = (a)^3 - (bi)^3 - 3(a)(bi)(a - bi) \\ & = a^3 - b^3(i)^3 - 3abi(a - bi) \\ & = a^3 - b^3(i)^2 \cdot i - 3a^2bi + \\ & \quad 3ab^2(i)^2 \\ & = a^3 - b^3(-1) \cdot i - 3a^2bi + \\ & \quad 3ab^2(-1) \\ & = a^3 - b^3i - 3a^2bi - 3ab^2 \end{aligned}$$

viii.  $(3 - \sqrt{-4})^{-3}$

$$\begin{aligned} \text{Solution:- } & (a - \sqrt{4}i)^{-3} \\ & = \frac{1}{(a - 2i)^3} \end{aligned}$$

$$\begin{aligned} & = \frac{1}{(3)^3 - (2i)^3 - 3(3)(2i)(3 - 2i)} \\ & = \frac{1}{27 - 2^3(i)^3 - 18i(3 - 2i)} \end{aligned}$$

$$\begin{aligned} & = \frac{1}{27 - 8(i)^2 \cdot i - 54i + 36i^2} \\ & = \frac{1}{27 - 8(-1) \cdot i - 54i + 36(-1)} \\ & = \frac{1}{27 + 8i - 54i - 36} \\ & = \frac{1}{-9 - 46i} \times \frac{-9 + 46i}{-9 + 46i} \\ & = \frac{(-9)^2 - (46i)^2}{-9 + 46i} \\ & = \frac{81 - 2116(i)^2}{-9 + 46i} \\ & = \frac{81 - 2116(-1)}{-9 + 46i} \\ & = \frac{81 + 2116}{-9 + 46i} \\ & = \frac{2197}{-9 + 46i} \\ & = \frac{-9}{2197} + \frac{46}{2197} \end{aligned}$$

A project of: [www.https://newsongoogle.com/](https://newsongoogle.com/)

Contact Us: bilalarticles1@gmail.com

